

Association between Dietary Intakes of Tea, Coffee, and Soft Drinks in Patients Undergoing Coronary Angiography with Coronary Artery Stenosis

Maryam Javadi^{1,2}, Atena Jamalzehi³, Hadis Gerami^{2,4,5}, Seyed Kianoosh Hosseini⁶, Mohammad Bagher Maljaei^{7,8,9}, Mina Eslami², Asma Bahreini⁹

¹Children Growth Research Center, Qazvin University of Medical Sciences, Qazvin, Iran, ²Department of Nutrition, School of Health, Qazvin University of Medical Sciences, Qazvin, Iran, ³Department of Nutrition, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran, ⁴Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran, ⁵Obesity and Eating Habits Research Center, Endocrinology and Metabolism Molecular Cellular Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran, ⁶Department of Cardiology, Tehran University of Medical Sciences Tehran, Iran, ⁷Students Research Committee, Iran University of Medical Sciences, Tehran, Iran, ⁸Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran, ⁹Isfahan Neuroscience Research Center, Alzahra Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: Coronary artery disease (CAD) is one of the major causes of mortality that is related to the nutritional habits and lifestyle. The aim of this study was to examine the association between tea, coffee, and soft drink consumption and coronary artery stenosis in patients undergoing coronary angiography.

Methods: Out of all the patients, 208 cases (101 Female) with 57.81 ± 12.18 (mean \pm SD) were assigned to participate in this cross-sectional study. In total, 168-items, semi-quantitative food frequency questionnaire collected for assessments of dietary intakes of black tea, coffee, caffeine, and soft drinks and record demographic and clinical questionnaire.

Results: There were negative association between arteries with stenosis of more than 50% number with dietary intakes of tea ($P = 0.011$, $r = -0.187$), coffee ($P = 0.069$, $r = -0.098$) intakes, and dietary caffeine intake ($P = 0.043$, $r = -0.118$). The high consumptions of soft drinks ($P = 0.005$, $r = 0.387$) were associated with an enhancement in arteries with stenosis of more than 50% number. In addition, dietary consumption of black tea have a negatively significant association with the history of previous angiography ($P = 0.044$, $r = -0.121$), the history of previous Stanton ($P = 0.035$, $r = -0.132$), and coronary artery bypass graft surgery nomination ($P = 0.008$, $r = -0.216$). Coffee consumption showed a significant negative relationship with engagement for coronary artery bypass graft surgery ($P = 0.004$, $r = -0.598$).

Conclusions: Dietary intakes of tea, coffee, and caffeine may have a negative relationship with CAD and cardio vascular diseases. Healthy dietary lifestyle is an important issue for the prevention of chronic diseases.

Keywords: Caffeine, carbonated beverages, coffee, coronary stenosis, tea